

Geometry and Topology Seminar
March 14, 2025, Lanzhou University



Spectrum and Strichartz estimate on locally symmetric spaces

Hong-Wei Zhang (Paderborn University)

Objects

(Continuous) Spectrum of
Laplace-Beltrami operator



Temperedness of $L^2(\Gamma \backslash G)$

Critical exponent of
discrete subgroup
(Poincaré series)

Application (original motivation): Strichartz inequality

Hyperbolic Space

Hyperbolic plane

$$\mathbb{H}^2 = \{z \in \mathbb{C} \mid \operatorname{Im} z > 0\}$$

(upper half-plane)

$$\mathbb{H}^2 = \operatorname{SL}(2, \mathbb{R})/\operatorname{SO}(2)$$

Real hyperbolic space

$$\mathbb{H}^n = \{x \in \mathbb{R} \times \mathbb{R}^n \mid -x_0^2 + x_1^2 + \cdots x_{n+1}^2 = -1, x_0 \geq 1\}$$

(hyperboloid)

$$\mathbb{H}^n = \operatorname{SO}_e(n+1, \mathbb{R})/\operatorname{SO}(n)$$

Hyperbolic Space

Hyperbolic plane

$$\mathbb{H}^2 = \{z \in \mathbb{C} \mid \operatorname{Im} z > 0\} \quad (\text{upper half-plane})$$

$$\mathbb{H}^2 = \operatorname{SL}(2, \mathbb{R})/\operatorname{SO}(2)$$

Real hyperbolic space

$$\mathbb{H}^n = \{x \in \mathbb{R} \times \mathbb{R}^n \mid -x_0^2 + x_1^2 + \cdots x_{n+1}^2 = -1, x_0 \geq 1\} \quad (\text{hyperboloid})$$

$$\mathbb{H}^n = \operatorname{SO}_e(n+1, \mathbb{R})/\operatorname{SO}(n)$$

Noncompact symmetric space of rank 1

$$\mathbb{H}^n = \mathbb{H}^n(\mathbb{R})$$

$$\mathbb{H}^n(\mathbb{C})$$

$$\mathbb{H}^n(\mathbb{H})$$

$$\mathbb{H}^2(\mathbb{O})$$


L^2 -Spectrum

- $-\Delta f = \lambda f$
- $\sigma(-\Delta) = \{\lambda \in \mathbb{C} \mid (-\Delta - \lambda)^{-1} : L^2 \rightarrow L^2 \text{ does not exist}\}$

L^2 -Spectrum

- $-\Delta f = \lambda f$
- $\sigma(-\Delta) = \{\lambda \in \mathbb{C} \mid (-\Delta - \lambda)^{-1} : L^2 \rightarrow L^2 \text{ does not exist}\}$


Euclidean setting




L^2 -Spectrum

- $-\Delta f = \lambda f$
- $\sigma(-\Delta) = \{\lambda \in \mathbb{C} \mid (-\Delta - \lambda)^{-1} : L^2 \rightarrow L^2 \text{ does not exist}\}$


Euclidean setting




Real Hyperbolic space



Hyperbolic Surface




Hyperbolic Surface




Modular curve $\text{SL}(2, \mathbb{Z}) \backslash \mathbb{H}^2$ (non-compact and finite area):



Hyperbolic Surface



Modular curve $\text{SL}(2, \mathbb{Z}) \backslash \mathbb{H}^2$ (non-compact and finite area):




i.e., there are

- infinitely many embedded eigenvalues
- no exceptional eigenvalues


(Selberg's 1/4 Conjecture for general Riemann surface)

Hyperbolic Surface



Thin group: $\Gamma \leq \text{SL}(2, \mathbb{R})$ s.t. $\text{Vol}(\Gamma \backslash \mathbb{H}^2) = \infty$


Hyperbolic Surface



Thin group: $\Gamma \leq \text{SL}(2, \mathbb{R})$ s.t. $\text{Vol}(\Gamma \backslash \mathbb{H}^2) = \infty$



Hyperbolic Surface




Thin group: $\Gamma \leq \text{SL}(2, \mathbb{R})$ s.t. $\text{Vol}(\Gamma \backslash \mathbb{H}^2) = \infty$



i.e., all eigenvalues are exceptional (finitely many)

Hyperbolic Surface



Thin group: $\Gamma \leq \mathrm{SL}(2, \mathbb{R})$ s.t. $\mathrm{Vol}(\Gamma \backslash \mathbb{H}^2) = \infty$




i.e., all eigenvalues are exceptional (finitely many)

Characterize $\lambda_0(\Gamma \backslash X) := \inf_{f \in \mathcal{C}_c^\infty(\Gamma \backslash X)} \frac{\int_{\Gamma \backslash X} \|\mathrm{grad} f\|^2 \, d\mathrm{vol}}{\int_{\Gamma \backslash X} \|f\|^2 \, d\mathrm{vol}} = \inf \sigma_c(-\Delta)$

Critical Exponent

$$\delta_\Gamma = \limsup_{R \rightarrow \infty} \frac{\log(\#\{\gamma \in \Gamma \mid d(e, \gamma e) \leq R\})}{R}$$




Critical Exponent

$$\delta_\Gamma = \limsup_{R \rightarrow \infty} \frac{\log(\#\{\gamma \in \Gamma \mid d(e, \gamma e) \leq R\})}{R}$$

$$\delta_\Gamma = \inf \left\{ s \in \mathbb{R} \mid \sum_{\gamma \in \Gamma} e^{-sd(e, \gamma e)} < \infty \right\}$$

Poincaré Series: $\sum_{\gamma \in \Gamma} e^{-sd(e, \gamma e)}$ $\begin{cases} < \infty & \text{if } s > \delta_\Gamma \\ = \infty & \text{if } s < \delta_\Gamma \end{cases}$


e.g. In \mathbb{H}^2 : $0 \leq \delta_\Gamma \leq 1$ In \mathbb{H}^n : $0 \leq \delta_\Gamma \leq n - 1$



Characterization in Dimension 2

$$\lambda_0(\Gamma \backslash \mathbb{H}^2) = \frac{1}{4}$$

$L^2(\Gamma \backslash X)$ is tempered


$$\delta(\Gamma) \leq \frac{1}{2}$$


Theorem [Elstrodt '73 *Math. Ann.*, Patterson '76 *Acta Math.*]

$$\lambda_0(\Gamma \backslash \mathbb{H}^2) = \begin{cases} \frac{1}{4} & \text{if } 0 \leq \delta_\Gamma \leq \frac{1}{2} \\ \frac{1}{4} - (\delta_\Gamma - \frac{1}{2})^2 & \text{if } \frac{1}{2} \leq \delta_\Gamma \leq 1 \end{cases}$$

Characterization on G/K of Rank 1

$$\lambda_0(\Gamma \backslash X) = \rho^2$$

$L^2(\Gamma \backslash X)$ is tempered


$$\delta(\Gamma) \leq \rho$$

Theorem [Elstrodt '73, Patterson '76, Sullivan '87 JDG, Corlette '90 *Invent. Math.*]

$$\lambda_0(\Gamma \backslash X) = \begin{cases} \rho^2 & \text{if } 0 \leq \delta_\Gamma \leq \rho \\ \rho^2 - (\delta_\Gamma - \rho)^2 & \text{if } \rho \leq \delta_\Gamma \leq 2\rho \end{cases}$$

where $\rho = \frac{n-1}{2}$ on $\mathbb{H}^n(\mathbb{R})$, n on $\mathbb{H}^n(\mathbb{C})$, $2n+1$ on $\mathbb{H}^n(\mathbb{H})$, 11 on $\mathbb{H}^2(\mathbb{O})$

Temperedness (柔曼性)

G connected semisimple Lie group \implies direct integrals:

$$L^2(\Gamma \backslash G) \cong \int_{\widehat{G}}^{\oplus} \mathcal{H}_{\pi} d\nu(\pi) \quad \text{and} \quad L^2(\Gamma \backslash X) \cong \int_{\widehat{G}_K}^{\oplus} (\mathcal{H}_{\pi})^K d\nu(\pi)$$

In rank 1, \widehat{G}_K consists of

- the unitary spherical principal series $\pi_{\pm\lambda}$ ($\lambda \in \mathbb{R}/\pm 1$)
- the trivial representation $\pi_{\pm i\rho} = 1$
- the complementary series $\pi_{\pm i\lambda}$ ($\lambda \in I$), where

$$I = \begin{cases} (0, \rho) & \text{if } X = \mathbb{H}^n(\mathbb{R}) \text{ or } \mathbb{H}^n(\mathbb{C}) \\ \left(0, \frac{m_{\alpha}}{2} + 1\right] & \text{if } X = \mathbb{H}^n(\mathbb{H}) \text{ or } \mathbb{H}^2(\mathbb{O}) \end{cases}$$

(no higher rank analogue)

Temperedness


$$\lambda_0(\Gamma \backslash X) = \rho^2 \quad \text{---} \quad L^2(\Gamma \backslash G) \text{ is tempered}$$
$$\delta(\Gamma) \leq \rho$$

```
graph TD; A["\lambda_0(\Gamma \backslash X) = \rho^2"] <--> B["L^2(\Gamma \backslash G) is tempered"]; C["\delta(\Gamma) \leq \rho"] --> A;
```

By definition

- $L^2(\Gamma \backslash G)$ is called tempered if \widehat{G}_K does not involve complementary series
- $-\Delta$ acts on $(\mathcal{H}_\pi)^K$ by multiplication by $\lambda^2 + \rho^2$

Question



② When $\Gamma \backslash X$ is of **higher rank** and infinite volume?

Noncompact (Riemannian) Symmetric Space

A noncompact symmetric space is a complete Riemannian manifold

- with **nonpositive** sectional curvature
- which is simply connected (Cartan-Hadamard manifold)

Noncompact (Riemannian) Symmetric Space

A noncompact symmetric space is a complete Riemannian manifold

- with nonpositive sectional curvature
- which is simply connected (Cartan-Hadamard manifold)
- with symmetric property
- which growths **exponentially** fast at infinity

Noncompact (Riemannian) Symmetric Space

A noncompact symmetric space is a complete Riemannian manifold

- with nonpositive sectional curvature
- which is simply connected (Cartan-Hadamard manifold)
- with symmetric property
- which grows exponentially fast at infinity
- which can be identified as a **homogeneous space** G/K

e.g. $\mathbb{H}^2 = \mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(2)$ and $\mathbb{H}^3 = \mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2)$

Locally Symmetric Space

Noncompact symmetric space

$$X = G/K$$

- G noncompact semisimple Lie group (connected, finite center)
- K maximal compact subgroup of G
- $\Gamma \leq G$: discrete and torsion-free subgroup of G
 - Γ is a lattice: $\text{Vol}(\Gamma \backslash X) < \infty$
 - Γ has infinite covolume: $\text{Vol}(\Gamma \backslash X) = \infty$

Weyl Chamber


Rank

Cartan subspace \mathfrak{a} : maximal connected, totally geodesic, flat sub-manifold of \mathbb{X}


$$\mathfrak{a} \approx \mathbb{R}^\ell \quad \text{and} \quad \ell = \dim \mathfrak{a} = \text{rank } G/K$$



A_1



$A_1 \times A_1$



A_2


Weyl Chamber

Rank

Cartan subspace \mathfrak{a} : maximal connected, totally geodesic, flat sub-manifold of \mathbb{X}

$$\mathfrak{a} \approx \mathbb{R}^\ell \quad \text{and} \quad \ell = \dim \mathfrak{a} = \text{rank } G/K$$

- Cartan decomposition $G = K(\exp \overline{\mathfrak{a}^+})K$




Weyl Chamber

Rank

Cartan subspace \mathfrak{a} : maximal connected, totally geodesic, flat sub-manifold of \mathbb{X}

$$\mathfrak{a} \approx \mathbb{R}^\ell \quad \text{and} \quad \ell = \dim \mathfrak{a} = \text{rank } G/K$$

- Cartan decomposition $G = K(\exp \overline{\mathfrak{a}^+})K$
- Cartan projection $\mu : G \longrightarrow \overline{\mathfrak{a}^+}$ such that
$$g \in Ke^{\mu(g)}K$$
- $d(e, \gamma e) = \|\mu(\gamma)\| \quad \forall \gamma \in \Gamma$



In higher rank: $\rho \in \mathfrak{a}^+$ is a **vector**, known as $\rho = \frac{1}{2} \sum_{\alpha \in \Sigma^+} m_\alpha \alpha$


Weyl Chamber

Rank

Cartan subspace \mathfrak{a} : maximal connected, totally geodesic, flat sub-manifold of \mathbb{X}

$$\mathfrak{a} \approx \mathbb{R}^\ell \quad \text{and} \quad \ell = \dim \mathfrak{a} = \text{rank } G/K$$

- Cartan decomposition $G = K(\exp \overline{\mathfrak{a}^+})K$
- Cartan projection $\mu : G \longrightarrow \overline{\mathfrak{a}^+}$ such that
$$g \in Ke^{\mu(g)}K$$
- $d(e, \gamma e) = \|\mu(\gamma)\| \quad \forall \gamma \in \Gamma$



In higher rank: $\rho \in \mathfrak{a}^+$ is a **vector**, known as $\rho = \frac{1}{2} \sum_{\alpha \in \Sigma^+} m_\alpha \alpha$

General Characterization

$$\lambda_0(\Gamma \setminus X) = \|\rho\|^2$$



$$\tilde{\delta}(\Gamma) \leq \|\rho\|$$

Theorem [Anker-Z. '22 *Geom. Dedicata*]

$$\lambda_0(\Gamma \setminus X) = \begin{cases} \|\rho\|^2 & \text{if } 0 \leq \tilde{\delta}_\Gamma \leq \|\rho\| \\ \|\rho\|^2 - (\tilde{\delta}_\Gamma - \|\rho\|)^2 & \text{if } \|\rho\| \leq \tilde{\delta}_\Gamma \leq 2\|\rho\| \end{cases}$$

Critical Exponent

- [Leuzinger '03]: Lower and upper bounds of $\lambda_0(\Gamma \backslash X)$ in terms of

$$\delta_\Gamma = \inf \left\{ s \in \mathbb{R} \mid \sum_{\gamma \in \Gamma} e^{-s \|\mu(\gamma)\|} < \infty \right\}$$

Critical Exponent

- [Leuzinger '03]: Lower and upper bounds of $\lambda_0(\Gamma \backslash X)$ in terms of


$$\delta_\Gamma = \inf \left\{ s \in \mathbb{R} \mid \sum_{\gamma \in \Gamma} e^{-s \|\mu(\gamma)\|} < \infty \right\}$$

- [Carron-Pedon '04, Anker-Z. '22]: Introduce the modified critical exponent

$$\tilde{\delta}_\Gamma = \inf \left\{ s \in \mathbb{R} \mid \sum_{\gamma \in \Gamma} e^{-\min\{s, \|\rho\|\} \langle \frac{\rho}{\|\rho\|}, \mu(\gamma) \rangle - \max\{0, s - \|\rho\|\} \|\mu(\gamma)\|} < \infty \right\}$$

- $0 \leq \delta_\Gamma \leq \tilde{\delta}_\Gamma \leq 2\|\rho\|$ and $\delta_\Gamma = \tilde{\delta}_\Gamma$ in rank 1

Critical Exponent



Temperedness

$$\lambda_0(\Gamma \backslash X) = \|\rho\|^2$$



$$\tilde{\delta}(\Gamma) \leq \|\rho\|$$

$L^2(\Gamma \backslash G)$ is tempered




$$\psi_\Gamma(H) \leq \rho(H) \quad \forall H \in \mathfrak{a}$$

Equivalence

- [Edwards-Oh '23 *Commun. Am. Math. Soc.*] : if Γ is Anosov
- [Lutsko-Weich-Wolf '24] : in general

Growth Indicator Function



$$\psi_{\Gamma}(H) = \|H\| \inf_{H \in \mathcal{C}} \inf \{s \in \mathbb{R} \mid \sum_{\gamma \in \Gamma, \mu(\gamma) \in \mathcal{C}} e^{-s\|\mu(\gamma)\|} < \infty\} \quad [\text{Quint '02 GAFA}]$$


$$\lambda_0(\Gamma \backslash X) = \|\rho\|^2$$

$L^2(\Gamma \backslash G)$ is tempered

$$\tilde{\delta}(\Gamma) \leq \|\rho\| \quad \xleftarrow{\text{red arrow}} \quad \psi_\Gamma(H) \leq \rho(H) \quad \forall H \in \mathfrak{a}$$

Theorem [Wolf-Z. '24 PAMS]

$$\tilde{\delta}_\Gamma = \begin{cases} \sup_{H \in \overline{\mathfrak{a}^+}} \psi_\Gamma(H) \cdot \frac{\|\rho\|}{\rho(H)} & \text{if } \psi_\Gamma \leq \rho \\ \sup_{H \in \overline{\mathfrak{a}^+}} \frac{\psi_\Gamma(H) - \rho(H)}{\|H\|} + \|\rho\| & \text{otherwise} \end{cases}$$




Theorem [Wolf-Z. '24 PAMS]

$$\lambda_0(\Gamma \backslash X) = \|\rho\|^2 - \max \left\{ 0, \sup_{H \in \overline{\mathfrak{a}_+}} \frac{\psi_\Gamma(H) - \rho(H)}{\|H\|} \right\}^2$$



Kunze-Stein Phenomenon

- Young's inequality: $L^p(G) * L^q(G) \subset L^r(G)$ $\frac{1}{p} + \frac{1}{q} = \frac{1}{r} + 1$
 $L^{\textcolor{red}{1}}(G) * L^2(G) \subset L^2(G)$



Kunze-Stein Phenomenon

- Young's inequality: $L^p(G) * L^q(G) \subset L^r(G)$ $\frac{1}{p} + \frac{1}{q} = \frac{1}{r} + 1$
 $L^1(G) * L^2(G) \subset L^2(G)$
- Kunze-Stein: $L^{\textcolor{red}{p}}(G) * L^2(G) \subset L^2(G)$ $1 \leq p < 2$



PDE Motivation

e.g. Free Schrödinger equation: $(i\partial_t + \Delta_x)u(t, x) = 0, \quad u(0, x) = f(x)$

whose solution is given by $u(t, x) = e^{it\Delta}f(x)$

Strichartz estimate

$$\|u\|_{L_t^p(\mathcal{I}, L_x^q(\mathcal{M}))} = \left(\int_{\mathcal{I}} dt \|u\|_{L^q(\mathcal{M})}^p \right)^{1/p} \lesssim \|f\|_{H^s(\mathcal{M})}$$

- for all **admissible pairs** (p, q)
- $s = 0$: without loss; $s > 0$: with loss of derivatives
- \mathcal{I} bounded: local-in-time; \mathcal{I} unbounded: global-in-time

Strichartz Estimates

e.g. Free Schrödinger equation: $(i\partial_t + \Delta_x)u(t, x) = 0, \quad u(0, x) = f(x)$

whose solution is given by $u(t, x) = e^{it\Delta}f(x)$

In \mathbb{R}^n [..., Strichartz '77 Duke, ..., Keel-Tao '98 AJM]

Global-in-time Strichartz inequality **without loss**

$$\|u\|_{L^p(\mathbb{R}, L^q(\mathbb{R}^n))} \lesssim \|f\|_{L^2(\mathbb{R}^n)}$$

holds for all **admissible** pairs (p, q) , i.e.,

$$\frac{2}{p} + \frac{n}{q} = \frac{n}{2} \quad p \geq 2 \quad (p, q) \neq (2, \infty)$$

Strichartz on manifolds

In \mathbb{R}^n

Global-in-time Strichartz holds without loss of derivatives

Compact manifold (M, g)

$$\|e^{it\Delta_g} u_0\|_{L_t^p(I, L_x^q(M))} = \left(\int_I dt \|e^{it\Delta_g} u_0\|_{L^q(M)}^p \right)^{1/p} \lesssim \|u_0\|_{H^s(M)}$$

- I is bounded

Strichartz on manifolds

In \mathbb{R}^n

Global-in-time Strichartz holds without loss of derivatives

Compact manifold (M, g)

$$\|e^{it\Delta_g} u_0\|_{L_t^p(I, L_x^q(M))} = \left(\int_I dt \|e^{it\Delta_g} u_0\|_{L^q(M)}^p \right)^{1/p} \lesssim \|u_0\|_{H^s(M)}$$

- I is bounded
- \mathbb{T}^n [Bourgain '93 *GAF*]: $s > \frac{n}{4} - \frac{1}{2}$
- M [Burq-Gérard-Tzvetkov '04 *AJM*]: $s = \frac{1}{p}$

Strichartz on manifolds

In \mathbb{R}^n

Global-in-time Strichartz holds without loss of derivatives

Compact manifold (M, g)

$$\|e^{it\Delta_g} u_0\|_{L_t^p(I, L_x^q(M))} = \left(\int_I dt \|e^{it\Delta_g} u_0\|_{L^q(M)}^p \right)^{1/p} \lesssim \|u_0\|_{H^s(M)}$$

- I is bounded
- \mathbb{T}^n [Bourgain '93 *GAF*]: $s > \frac{n}{4} - \frac{1}{2}$
- M [Burq-Gérard-Tzvetkov '04 *AJM*]: $s = \frac{1}{p}$


Question: on which manifolds does Strichartz hold without any loss ?

Strichartz on G/K

Global-in-time Strichartz estimate

$$\|e^{it\Delta} u_0\|_{L_t^p(\mathbb{R}, L_x^q(G/K))} \lesssim \|u_0\|_{L^2(G/K)}$$

holds without any loss of derivatives for all (p, q) admissible:



On Locally Symmetric Space

Strichartz estimate

Global-in-time Strichartz inequality holds without lossing any derivatives for the large X -admissible set if the following conditions are met:

- X has rank 1
- Γ is convex cocompact
- $\delta_\Gamma < \rho$

[Burq-Guillarmou-Hassell '10 GAFA, Fotiadis-Mandouvalos-Marias '18 Math. Ann.]




Figure: Convex hull of the limit set Λ_Γ in \mathbb{H}^2

$$\begin{array}{c} \Gamma \text{ convex cocompact} \\ \Updownarrow \\ \Gamma \backslash \text{Conv}(\Lambda_\Gamma) \text{ compact} \end{array}$$

On Locally Symmetric Space

Strichartz estimate

Global-in-time Strichartz inequality holds without lossing any derivatives for the large X -admissible set if the following conditions are met:

- X has rank 1
- Γ is convex cocompact
- $\delta_\Gamma < \rho$

[Burq-Guillarmou-Hassell '10 *GAFA*, Fotiadis-Mandouvalos-Marias '18 *Math. Ann.*]

Remark.

- δ_Γ small enough \implies Γ is convex cocompact [Liu-Wang '23 *GT*]
- $\delta_\Gamma < \rho$ \implies temperedness \implies Kunze-Stein [Z. '20 *JGA*]