

Analysis Disguised as Topology: Poincaré Duality and the Hodge Star Map

Edward Bryden

University of Antwerp

November 10, 2025

Table of Contents

1 Hodge Theory

2 Real Homology

3 Poincaré Duality

4 Hodge Star Operator

Adjoint of d

Recall

Let (M, g) be a smooth closed m -dimensional Riemannian manifold. Recall that we gave the adjoint of d the special name δ ; it is a map $\delta : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$.

Adjoint of d

Lemma

Let (M, g) be a smooth closed m -dimensional Riemannian manifold, and let δ be the adjoint of the exterior derivative d . Then, $\delta^2 = 0$.

Proof.

By definition, for all $\omega \in \Omega^{k-2}(M)$ and $\eta \in \Omega^k(M)$ we have

$$\int_M g(\omega, \delta^2 \eta) = \int_M g(d^2 \omega, \eta) = 0. \quad (1)$$



The Hodge Laplacian

Definition

Let (M, g) be a smooth closed m -dimensional Riemannian manifold. Then, we define the Hodge-Laplacian $\Delta : \Omega^k(M) \rightarrow \Omega^k(M)$ by

$$\Delta\sigma = (d + \delta)^2\sigma = (d\delta + \delta d)\sigma. \quad (2)$$

If $\Delta\sigma = 0$, then we call σ an harmonic form.

Lemma

Let (M, g) be a smooth closed m -dimensional Riemannian manifold. Then, every element $w \in H^k(M)$ has a unique harmonic representative $\omega \in \Omega^k(M)$. Furthermore, we have that

$$\int_M g(\omega, \omega) = \min \left\{ \int_M g(\sigma, \sigma) : \sigma \in w \right\}. \quad (3)$$

Consequences for the integral cohomology lattice?

Recall

Recall that we defined $H^1(M; \mathbb{Z})_{\mathbb{R}}$ to be those elements of $H^1_{dR}(M)$ whose representatives ω satisfy

$$\int_{\gamma} \omega \in \mathbb{Z} \forall \gamma \in \pi_1(M). \quad (4)$$

Observation

Since the topological properties of $H^1(M; \mathbb{Z})_{\mathbb{R}}$ are tied to $\pi_1(M)$, one may expect that the geometric properties of $H^1(M; \mathbb{Z})_{\mathbb{R}}$ are linked to the geometric properties of $\pi_1(M)$.


One-cycles

Definition

Let M be a smooth manifold, and let Δ^1 denote the 1-dimensional simplex $[0, 1]$. Recall that given $\sigma : [0, 1] \rightarrow M$ we set $\partial\sigma = \sigma(1) - \sigma(0)$. A real one cycle is a finite sum of $\sigma_i : [0, 1] \rightarrow M$ with coefficients $r_i \in \mathbb{R}$ such that

$$\partial \sum_i r_i \sigma_i = \sum_i r_i \partial \sigma_i = 0. \quad (5)$$

Sketch



Real k -cycles

Definition

Let M be a smooth manifold, and let Δ^k denote the k -dimensional simplex. A real k -cycle is a finite sum of $\sigma_i : \Delta^k \rightarrow M$ with coefficients $r_i \in \mathbb{R}$ such that


$$\partial \sum_i r_i \sigma_i = \sum_i r_i \partial \sigma_i = 0. \quad (6)$$

Definition

Let M be a smooth manifold, and let c_1 and c_2 be two real one-cycles. We say that they are homologous if there is a real two-cycle σ such that

$$\partial\sigma = c_2 - c_1. \quad (7)$$

Sketch



Definition

Let M be a smooth manifold. Two real k -cycles c_1 and c_2 are said to be homologous if and only if there exists a $k + 1$ cycle σ such that $\partial\sigma = c_2 - c_1$. We denote the collection of all k -real-homology classes of M by $H_k(M; \mathbb{R})$

Homology-Cohomology Pairing

Definition

Let M be a closed smooth manifold. Given a map $f : \Delta^k \rightarrow M$ we get a corresponding map $\Omega^k(M) \rightarrow \mathbb{R}$ given by

$$\omega \mapsto \int_{\Delta^k} f^* \omega. \quad (8)$$

Lemma

This map descends to a map $I : H_k(M; \mathbb{R}) \times H_{dR}^k(M) \rightarrow \mathbb{R}$.

Proof.

This is an application of Stoke's theorem to the relevant definitions. □

The Pairing is Non-degenerate

Theorem

Let M be a closed smooth oriented manifold. Then, the map $I : H_k(M; \mathbb{R}) \times H_{dR}^k(M) \rightarrow \mathbb{R}$ is non-degenerate in the following sense.

- 1 For $a \in H_k(M; \mathbb{R})$ we have $I(a, w) = 0$ for all $w \in H_{dR}^k(M)$ if and only if $a = 0$.
- 2 For $w \in H_{dR}^k(M)$ we have $I(a, w) = 0$ for all $a \in H_k(M; \mathbb{R})$ if and only if $w = 0$.

Cohomology-Cohomology Pairing

Definition

Let M be a smooth closed oriented m -dimensional manifold. We get a pairing $I_\wedge : \Omega^k(M) \times \Omega^{m-k}(M) \rightarrow \mathbb{R}$ as follows:

$$I_\wedge(\eta, \omega) = \int_M \eta \wedge \omega. \quad (9)$$

Lemma


The pairing I_\wedge descends to $I_\wedge : H_{dR}^k(M) \times H_{dR}^{m-k}(M)$. Here it is non-degenerate.

Definition


Let M be a smooth closed m -dimensional manifold, and consider $a \in H_k(M; \mathbb{R})$. The Poincaré Dual of a is the unique element $\text{PD}(a) \in H_{dR}^{m-k}(M; \mathbb{R})$ such that for all $h \in H_{dR}^k(M)$ we have

$$I(a, h) = I_{\wedge}(\text{PD}(a), h). \quad (10)$$

Sketch



Sketch



Definition

Let M be a smooth closed oriented m -dimensional manifold, and consider $w \in H_{dR}^k(M)$. The Poincaré Dual of w is the unique element $\text{PD}(w)$ in $H_{m-k}(M; \mathbb{R})$ such that for all $h \in H_{m-k}(M; \mathbb{R})$ such that

$$I(\text{PD}(w), h) = I_{\wedge}(w, h). \quad (11)$$

Interior Product

Definition

Let (M, g) be a smooth m -dimensional Riemannian manifold, let $k \leq m$, let $p \in M$, and $\eta \in \bigwedge_{i=1}^k T_p^* M$. Then, for each $l \leq m - k$ we get a map $\psi \mapsto \eta \wedge \psi$ from $\bigwedge_{i=1}^l T_p^* M$ to $\bigwedge_{i=1}^{l+k} T_p^* M$. Define $\eta_{\lrcorner} : \bigwedge_{i=1}^{l+k} T_p^* M \rightarrow \bigwedge_{i=1}^l T_p^* M$ to be its adjoint: for all $\omega \in \bigwedge_{i=1}^{l+k} T_p^* M$ and all $\sigma \in \bigwedge_{i=1}^l T_p^* M$ we have

$$g(\eta_{\lrcorner}(\omega), \sigma) = g(\omega, \eta \wedge \sigma). \quad (12)$$

Hodge Star Map

Definition

Let (M, g) be a smooth oriented m – *dimensional* Riemannian manifold, and let vol_g denote its volume form. For any $k \leq m$ we define a map $\star : \Omega^k(M) \rightarrow \Omega^{m-k}(M)$ as follows. For $\sigma \in \Omega^k(M)$ for each $p \in M$ we set

$$(\star\sigma)_p = (\sigma_p)_{\lrcorner}(\text{vol}_g)_p. \quad (13)$$

Lemma

Let (M, g) be a smooth closed oriented m -dimensional manifold. Then for each $p \in M$ and all $\omega, \sigma \in \bigwedge_{i=1}^k T_p^* M$ we have

$$g(\omega, \sigma) \cdot (\text{vol}_g)_p = \omega \wedge \star \sigma. \quad (14)$$

By integrating over M , we get

$$\int_M g(\omega, \sigma) \text{vol}_g = \int_M \omega \wedge \star \sigma. \quad (15)$$

Proof.

Let $\{E_i\}_{i=1}^m$ be an orthonormal basis for $T_p M$, and let $\{\Theta_i\}_{i=1}^m$ be the dual basis for $T_p^* M$. Consider the element $\sigma = \Theta_1 \wedge \cdots \wedge \Theta_k$, and observe that $\sigma \lrcorner \text{vol}_g = \Theta_{k+1} \wedge \cdots \wedge \Theta_m$. This shows that for any ω we have

$$g(\omega, \Theta_1 \wedge \cdots \wedge \Theta_k) \cdot (\text{vol}_g)_p = \omega \wedge \Theta_{k+1} \wedge \cdots \wedge \Theta_m. \quad (16)$$

Similar arguments work for all of the elements Θ_J for $J \in J(k, m)$. □

Corollary

The map \star satisfies $\star^2 = (-1)^{k(m-k)} \text{Id}$.

Corollary

The map \star is an isometry.

Proof.

$$g(\star\omega, \star\omega)\text{vol}_g = \star\omega \wedge \star^2\omega \quad (17)$$

$$= (-1)^{k(m-k)} \star\omega \wedge \omega \quad (18)$$

$$= (-1)^{k(m-k)} (-1)^{k(m-k)} \omega \wedge \star\omega \quad (19)$$

$$= g(\omega, \omega)\text{vol}_g. \quad (20)$$



Length

Definition

Let (M, g) be a smooth Riemannian manifold, and let $\sigma : [0, 1] \rightarrow M$ be a curve. Then, the length of the curve, denoted by $\text{vol}_1([0, 1], \sigma^*g)$ is given by

$$\int_0^1 \sqrt{g(\dot{\sigma}, \dot{\sigma})} dt. \quad (21)$$

Observation

The integrand is the area form on $[0, 1]$ induced by the metric σ^*g on $[0, 1]$.

Definition

Let (M, g) be a smooth closed m -dimensional Riemannian manifold, and let Δ^k denote the k -simplex. Then, for $f : \Delta^k \rightarrow M$ we define $\text{vol}_k(\Delta^k, f^*g)$ to be

$$\text{vol}_k(\Delta^k, f^*g) = \int_{\Delta^k} \text{vol}_{f^*g} \quad (22)$$

Warning

The fully accurate definition is a bit more subtle

Definition

Let (M, g) be a smooth closed m -dimensional Riemannian manifold. Given a class $w \in H_k(M; \mathbb{R})$ we define its k -mass to be

$$\|w\|_k = \inf \left\{ \sum_i |r_i| \text{vol}_k(\Delta^k, \sigma_i^* g) : \sum_i r_i \sigma_i \in w \right\} \quad (23)$$

Lemma (Hebda)

Let (M, g) be a smooth closed oriented Riemannian manifold. For $a \in H_{dR}^{m-1}(M)$ and $\text{PD}(a) \in H_1(M; \mathbb{R})$ we have

$$\|\text{PD}(a)\|_1 \leq \text{Vol}_g(M)^{\frac{1}{2}} C(m, 1) \inf \left\{ \left(\int_M g(\omega, \omega) \right)^{\frac{1}{2}} : \omega \in a \right\} \quad (24)$$

Proof.

Let ω be the harmonic form representing a . Then, for all closed 1-forms ϕ we have

$$I(\text{PD}(a), \phi) = I_{\wedge}(\omega, \phi) = \int_M \omega \wedge \phi. \quad (25)$$

This RHS is equal to

$$\pm \int_M g(\star \omega, \phi). \quad (26)$$

Proof.

Taking absolute values, we get

$$|I(\text{PD}(a), \phi)| \leq \int_M |\star \omega|_g |\phi| \quad (27)$$

$$\leq \|\phi\|_{L^\infty} \int_M |\star \omega| \quad (28)$$

$$\leq \|\phi\|_{L^\infty} \text{vol}_g(M)^{\frac{1}{2}} \left(\int_M g(\star \omega, \star \omega) \right)^{\frac{1}{2}}. \quad (29)$$

$$= \|\phi\|_{L^\infty} \text{vol}_g(M)^{\frac{1}{2}} \left(\int_M g(\omega, \omega) \right)^{\frac{1}{2}}. \quad (30)$$

Proof.

On the otherhand, by the properties of mass, we have

$$\|\mathrm{PD}(a)\|_1 \leq C(m, 1) \sup \{ |I(\mathrm{PD}(a), \phi)| : \|\phi\|_{L^\infty} \leq 1 \}. \quad (31)$$

